

COMPRESSED RANDOM VARIABLES IN THE GRAPH W^* -PROBABILITY SPACES

ILWOO CHO

ABSTRACT. In [16] and [17], we observed the amalgamated free probability theory on the graph W^* -probability space $(W^*(G), E)$ over the diagonal subalgebra D_G . In [18], we consider the diagonal compressed random variables in $(W^*(G), E)$ and observed the amalgamated freeness in the W^* -probability space $(W^*(G), E)$ over the D_G . In particular, for $V_N = \{v_0\} \subset V(G)$, we could construct the (scalar-valued) tracial W^* -probability space, so-called the vertex compressed graph W^* -probability space. In this paper, we will consider the off-diagonal compressed random variables in $(W^*(G), E)$. After fixing $v_1 \neq v_2$ in $V(G)$, we define the (v_1, v_2) -off-diagonal compressed random variable of $a \in (W^*(G), E)$, $L_{v_1} a L_{v_2}$. We will consider the free probability data on such elements in $(W^*(G), E)$. Also, we will consider the compressed random variables PaP for the D_G -valued random variable $a \in (W^*(G), E)$, compressed by the projection $P = L_{v_1} + L_{v_2} + \dots + L_{v_N} \in D_G$. To observe the free probability data for such compressed random variable, we use the diagonal-compression and off-diagonal compression. We can figure out that only the diagonal-compression affects the compressed free probability on $W^*(G)$. In fact, D_G -valued moment series and R-transform of the compressed random variable of a are same as those of diagonal compressed random of it.

In [16], we constructed the graph W^* -probability spaces. The graph W^* -probability theory is one of the good example of Speicher's combinatorial free probability theory with amalgamation. In [16], we observed how to compute the moment and cumulant of an arbitrary random variables in the graph W^* -probability space and the freeness on it with respect to the given conditional expectation. Also, in [17], we consider certain special random variables of the graph W^* -probability space, for example, semicircular elements, even elements and R-diagonal elements. This shows that the graph W^* -probability spaces contain the rich free probabilistic objects. Roughly speaking, graph W^* -algebras are W^* -topology closed version of free semigroupoid algebras defined and observed by Kribs and Power in [10].

Throughout this paper, let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G , in the sense of Kribs and Power. i.e., it is a collection of all vertices of the graph G as units and all admissible finite paths, under the admissibility. As a set, the free semigroupoid $\mathbb{F}^+(G)$ can be decomposed by

$$\mathbb{F}^+(G) = V(G) \cup FP(G),$$

Key words and phrases. Graph W^* -Probability Spaces over Diagonal Subalgebras, Diagonal-Compressed Random Variables, Off-Diagonal Compressed Random Variables. Compressed Random Variables.

where $V(G)$ is the vertex set of the graph G and $FP(G)$ is the set of all admissible finite paths. Trivially the edge set $E(G)$ of the graph G is properly contained in $FP(G)$, since all edges of the graph can be regarded as finite paths with their length 1. We define a graph W^* -algebra of G by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w,$$

where L_w and L_w^* are creation operators and annihilation operators on the generalized Fock space $H_G = l^2(\mathbb{F}^+(G))$ induced by the given graph G , respectively. Notice that the creation operators induced by vertices are projections and the creation operators induced by finite paths are partial isometries. We can define the W^* -subalgebra D_G of $W^*(G)$, which is called the diagonal subalgebra by

$$D_G \stackrel{def}{=} \overline{\mathbb{C}[\{L_v : v \in V(G)\}]}^w.$$

Then each element a in the graph W^* -algebra $W^*(G)$ is expressed by

$$a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}, \text{ for } p_w \in \mathbb{C},$$

where $\mathbb{F}^+(G : a)$ is a support of the element a , as a subset of the free semi-groupoid $\mathbb{F}^+(G)$. The above expression of the random variable a is said to be the Fourier expansion of a . Since $\mathbb{F}^+(G)$ is decomposed by the disjoint subsets $V(G)$ and $FP(G)$, the support $\mathbb{F}^+(G : a)$ of a is also decomposed by the following disjoint subsets,

$$V(G : a) = \mathbb{F}^+(G : a) \cap V(G)$$

and

$$FP(G : a) = \mathbb{F}^+(G : a) \cap FP(G).$$

Thus the operator a can be re-expressed by

$$a = \sum_{v \in V(G:a)} p_v L_v + \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

Notice that if $V(G : a) \neq \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v$ is contained in the diagonal subalgebra D_G . Thus we have the canonical conditional expectation $E : W^*(G) \rightarrow D_G$, defined by

$$E(a) = \sum_{v \in V(G:a)} p_v L_v,$$

for all $a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1,*\}} p_w L_w^{u_w}$ in $W^*(G)$. Then the algebraic pair $(W^*(G), E)$ is a W^* -probability space with amalgamation over D_G (See [16]). It is easy to check that the conditional expectation E is faithful in the sense that if $E(a^*a) = 0_{D_G}$, for $a \in W^*(G)$, then $a = 0_{D_G}$.

For the fixed operator $a \in W^*(G)$, the support $\mathbb{F}^+(G : a)$ of the operator a is again decomposed by

$$\mathbb{F}^+(G : a) = V(G : a) \cup FP_*(G : a) \cup FP_*^c(G : a),$$

with the decomposition of $FP(G : a)$,

$$FP(G : a) = FP_*(G : a) \cup FP_*^c(G : a),$$

where

$$FP_*(G : a) = \{w \in FP(G : a) : \text{both } L_w \text{ and } L_w^* \text{ are summands of } a\}$$

and

$$FP_*^c(G : a) = FP(G : a) \setminus FP_*(G : a).$$

The above new expression plays a key role to find the D_G -valued moments of the random variable a . In fact, the summands $p_v L_v$'s and $p_w L_w + p_w^* L_w^*$, for $v \in V(G : a)$ and $w \in FP_*(G : a)$ act for the computation of D_G -valued moments of a . By using the above partition of the support of a random variable, we can compute the D_G -valued moments and D_G -valued cumulants of it via the lattice path model LP_n and the lattice path model LP_n^* satisfying the $*$ -axis-property. At a first glance, the computations of D_G -valued moments and cumulants look so abstract and hence it looks useless. However, these computations, in particular the computation of D_G -valued cumulants, provides us how to figure out the D_G -freeness of random variables by making us compute the mixed cumulants. As applications, in the final chapter, we can compute the moment and cumulant of the operator that is the sum of N -free semicircular elements with their covariance 2.

Based on the D_G -cumulant computation, we can characterize the D_G -freeness of generators of $W^*(G)$, by the so-called diagram-distinctness on the graph G . i.e., the random variables L_{w_1} and L_{w_2} are free over D_G if and only if w_1 and w_2 are diagram-distinct the sense that w_1 and w_2 have different diagrams on the graph G . Also, we could find the necessary condition for the D_G -freeness of two arbitrary random variables a and b . i.e., if the supports $\mathbb{F}^+(G : a)$ and $\mathbb{F}^+(G : b)$ are diagram-distinct, in the sense that w_1 and w_2 are diagram distinct for all pairs $(w_1, w_2) \in \mathbb{F}^+(G : a) \times \mathbb{F}^+(G : b)$, then the random variables a and b are free over D_G .

In [17], we considered some special D_G -valued random variables in a graph W^* -probability space $(W^*(G), E)$. The those random variables are the basic objects to study Free Probability Theory. We can conclude that

- (i) if l is a loop, then $L_l + L_l^*$ is D_G -semicircular.

(ii) if w is a finite path, then $L_w + L_w^*$ is D_G -even.

(iii) if w is a finite path, then L_w and L_w^* are D_G -valued R-diagonal.

In [18], we observed the diagonal compressed random variables in the graph W^* -probability space $(W^*(G), E)$. Let $v_1, \dots, v_N \in V(G)$ and let a be a D_G -valued random variable in $(W^*(G), E)$. Define the diagonal compressed random variable of a by $V = \{v_1, \dots, v_N\}$ by

$$C_V(a) = L_{v_1}aL_{v_1} + \dots + L_{v_N}aL_{v_N}.$$

Notice that if $v \in V(G)$, then $L_v a L_v$ is the compressed random variable by L_v and the compressed random variable has its support contained in $\{v\} \cup \text{loop}_v(G)$, where $\text{loop}_v(G) = \{l \in \text{loop}(G) : l = vlv\}$.

The main purpose of this paper is to show that the compressed random variable $P_V a P_V$ of a projection $P_V = \sum_{j=1}^N L_{v_j}$, where $V = \{v_1, \dots, v_N\}$, has the same free probabilistic information with the diagonal compressed random variable $C_V(a)$. i.e., the compressed random variables $P_V a P_V$ and $C_V(a)$ have the same D_G -valued moments and cumulants.

1. GRAPH W^* -PROBABILITY THEORY

Let G be a countable directed graph and let $\mathbb{F}^+(G)$ be the free semigroupoid of G . i.e., the set $\mathbb{F}^+(G)$ is the collection of all vertices as units and all admissible finite paths of G . Let w be a finite path with its source $s(w) = x$ and its range $r(w) = y$, where $x, y \in V(G)$. Then sometimes we will denote w by $w = xwy$ to express the source and the range of w . We can define the graph Hilbert space H_G by the Hilbert space $l^2(\mathbb{F}^+(G))$ generated by the elements in the free semigroupoid $\mathbb{F}^+(G)$. i.e., this Hilbert space has its Hilbert basis $\mathcal{B} = \{\xi_w : w \in \mathbb{F}^+(G)\}$. Suppose that $w = e_1 \dots e_k \in FP(G)$ is a finite path with $e_1, \dots, e_k \in E(G)$. Then we can regard ξ_w as $\xi_{e_1} \otimes \dots \otimes \xi_{e_k}$. So, in [10], Kribs and Power called this graph Hilbert space the generalized Fock space. Throughout this paper, we will call H_G the graph Hilbert space to emphasize that this Hilbert space is induced by the graph.

Define the creation operator L_w , for $w \in \mathbb{F}^+(G)$, by the multiplication operator by ξ_w on H_G . Then the creation operator L on H_G satisfies that

(i) $L_w = L_{xwy} = L_x L_w L_y$, for $w = xwy$ with $x, y \in V(G)$.

$$(ii) \ L_{w_1} L_{w_2} = \begin{cases} L_{w_1 w_2} & \text{if } w_1 w_2 \in \mathbb{F}^+(G) \\ 0 & \text{if } w_1 w_2 \notin \mathbb{F}^+(G), \end{cases}$$

for all $w_1, w_2 \in \mathbb{F}^+(G)$.

Now, define the annihilation operator L_w^* , for $w \in \mathbb{F}^+(G)$ by

$$L_w^* \xi_{w'} \stackrel{def}{=} \begin{cases} \xi_h & \text{if } w' = wh \in \mathbb{F}^+(G) \\ 0 & \text{otherwise.} \end{cases}$$

The above definition is gotten by the following observation ;

$$\begin{aligned} \langle L_w \xi_h, \xi_{wh} \rangle &= \langle \xi_{wh}, \xi_{wh} \rangle \\ &= 1 = \langle \xi_h, \xi_h \rangle \\ &= \langle \xi_h, L_w^* \xi_{wh} \rangle, \end{aligned}$$

where $\langle, \rangle$ is the inner product on the graph Hilbert space H_G . Of course, in the above formula we need the admissibility of w and h in $\mathbb{F}^+(G)$. However, even though w and h are not admissible (i.e., $wh \notin \mathbb{F}^+(G)$), by the definition of L_w^* , we have that

$$\begin{aligned} \langle L_w \xi_h, \xi_h \rangle &= \langle 0, \xi_h \rangle \\ &= 0 = \langle \xi_h, 0 \rangle \\ &= \langle \xi_h, L_w^* \xi_h \rangle. \end{aligned}$$

Notice that the creation operator L and the annihilation operator L^* satisfy that

$$(1.1) \quad L_w^* L_w = L_y \quad \text{and} \quad L_w L_w^* = L_x, \quad \text{for all } w = xwy \in \mathbb{F}^+(G),$$

under the weak topology, where $x, y \in V(G)$. Remark that if we consider the von Neumann algebra $W^*(\{L_w\})$ generated by L_w and L_w^* in $B(H_G)$, then the projections L_y and L_x are Murray-von Neumann equivalent, because there exists a partial isometry L_w satisfying the relation (1.1). Indeed, if $w = xwy$ in $\mathbb{F}^+(G)$, with $x, y \in V(G)$, then under the weak topology we have that

$$(1.2) \quad L_w L_w^* L_w = L_w \quad \text{and} \quad L_w^* L_w L_w^* = L_w^*.$$

So, the creation operator L_w is a partial isometry in $W^*(\{L_w\})$ in $B(H_G)$. Assume now that $v \in V(G)$. Then we can regard v as $v = vvv$. So,

$$(1.3) \quad L_v^* L_v = L_v = L_v L_v^* = L_v^*.$$

This relation shows that L_v is a projection in $B(H_G)$ for all $v \in V(G)$.

Define the **graph W^* -algebra** $W^*(G)$ by

$$W^*(G) \stackrel{def}{=} \overline{\mathbb{C}[\{L_w, L_w^* : w \in \mathbb{F}^+(G)\}]}^w.$$

Then all generators are either partial isometries or projections, by (1.2) and (1.3). So, this graph W^* -algebra contains a rich structure, as a von Neumann algebra. (This construction can be the generalization of that of group von Neumann algebra.) Naturally, we can define a von Neumann subalgebra $D_G \subset W^*(G)$ generated by all projections L_v , $v \in V(G)$. i.e.

$$D_G \stackrel{def}{=} W^*(\{L_v : v \in V(G)\}).$$

We call this subalgebra the **diagonal subalgebra** of $W^*(G)$. Notice that $D_G = \Delta_{|G|} \subset M_{|G|}(\mathbb{C})$, where $\Delta_{|G|}$ is the subalgebra of $M_{|G|}(\mathbb{C})$ generated by all diagonal matrices. Also, notice that $1_{D_G} = \sum_{v \in V(G)} L_v = 1_{W^*(G)}$.

If $a \in W^*(G)$ is an operator, then it has the following decomposition which is called the Fourier expansion of a ;

$$(1.4) \quad a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w},$$

where $p_w \in \mathbb{C}$ and $\mathbb{F}^+(G:a)$ is the support of a defined by

$$\mathbb{F}^+(G:a) = \{w \in \mathbb{F}^+(G) : p_w \neq 0\}.$$

Remark that the free semigroupoid $\mathbb{F}^+(G)$ has its partition $\{V(G), FP(G)\}$, as a set. i.e.,

$$\mathbb{F}^+(G) = V(G) \cup FP(G) \quad \text{and} \quad V(G) \cap FP(G) = \emptyset.$$

So, the support of a is also partitioned by

$$\mathbb{F}^+(G:a) = V(G:a) \cup FP(G:a),$$

where

$$V(G:a) \stackrel{def}{=} V(G) \cap \mathbb{F}^+(G:a)$$

and

$$FP(G:a) \stackrel{def}{=} FP(G) \cap \mathbb{F}^+(G:a).$$

So, the above Fourier expansion (1.4) of the random variable a can be re-expressed by

$$(1.5) \quad a = \sum_{v \in V(G:a)} p_v L_v + \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

We can easily see that if $V(G : a) \neq \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v$ is contained in the diagonal subalgebra D_G . Also, if $V(G : a) = \emptyset$, then $\sum_{v \in V(G:a)} p_v L_v = 0_{D_G}$. So, we can define the following canonical conditional expectation $E : W^*(G) \rightarrow D_G$ by

$$(1.6) \quad E(a) = E \left(\sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1,*\}} p_w L_w^{u_w} \right) \stackrel{def}{=} \sum_{v \in V(G:a)} p_v L_v,$$

for all $a \in W^*(G)$. Indeed, E is a well-determined conditional expectation.

Definition 1.1. Let G be a countable directed graph and let $W^*(G)$ be the graph W^* -algebra induced by G . Let $E : W^*(G) \rightarrow D_G$ be the conditional expectation defined above. Then we say that the algebraic pair $(W^*(G), E)$ is the graph W^* -probability space over the diagonal subalgebra D_G . By the very definition, it is one of the W^* -probability space with amalgamation over D_G . All elements in $(W^*(G), E)$ are called D_G -valued random variables.

We have a graph W^* -probability space $(W^*(G), E)$ over its diagonal subalgebra D_G . We will define the following free probability data of D_G -valued random variables.

Definition 1.2. Let $W^*(G)$ be the graph W^* -algebra induced by G and let $a \in W^*(G)$. Define the n -th (D_G -valued) moment of a by

$$E(d_1 a d_2 a \dots d_n a), \text{ for all } n \in \mathbb{N},$$

where $d_1, \dots, d_n \in D_G$. Also, define the n -th (D_G -valued) cumulant of a by

$$k_n(d_1 a, d_2 a, \dots, d_n a) = C^{(n)}(d_1 a \otimes d_2 a \otimes \dots \otimes d_n a),$$

for all $n \in \mathbb{N}$, and for $d_1, \dots, d_n \in D_G$, where $\widehat{C} = (C^{(n)})_{n=1}^\infty \in I^c(W^*(G), D_G)$ is the cumulant multiplicative bimodule map induced by the conditional expectation E , in the sense of Speicher. We define the n -th trivial moment of a and the n -th trivial cumulant of a by

$$E(a^n) \quad \text{and} \quad k_n \left(\underbrace{a, a, \dots, a}_{n\text{-times}} \right) = C^{(n)}(a \otimes a \otimes \dots \otimes a),$$

respectively, for all $n \in \mathbb{N}$.

To compute the D_G -valued moments and cumulants of the D_G -valued random variable a , we need to introduce the following new definition ;

Definition 1.3. Let $(W^*(G), E)$ be a graph W^* -probability space over D_G and let $a \in (W^*(G), E)$ be a random variable. Define the subset $FP_*(G : a)$ in $FP(G : a)$ by

$$FP_*(G : a) \stackrel{def}{=} \{w \in \mathbb{F}^+(G : a) : \text{both } L_w \text{ and } L_w^* \text{ are summands of } a\}.$$

$$\text{And let } FP_*^c(G : a) \stackrel{def}{=} FP(G : a) \setminus FP_*(G : a).$$

We already observed that if $a \in (W^*(G), E)$ is a D_G -valued random variable, then a has its Fourier expansion $a_d + a_0$, where

$$a_d = \sum_{v \in V(G:a)} p_v L_v$$

and

$$a_0 = \sum_{w \in FP(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

By the previous definition, the set $FP(G : a)$ is partitioned by

$$FP(G : a) = FP_*(G : a) \cup FP_*^c(G : a),$$

for the fixed random variable a in $(W^*(G), E)$. So, the summand a_0 , in the Fourier expansion of $a = a_d + a_0$, has the following decomposition ;

$$a_0 = a_{(*)} + a_{(non-*)},$$

where

$$a_{(*)} = \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^*} L_l^*)$$

and

$$a_{(non-*)} = \sum_{w \in FP_*^c(G:a), u_w \in \{1, *\}} p_w L_w^{u_w},$$

where p_{l^*} is the coefficient of L_l^* depending on $l \in FP_*(G : a)$.

1.1. D_G -Moments and D_G -Cumulants of Random Variables.

Throughout this chapter, let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . In this chapter, we will compute the D_G -valued moments and the D_G -valued cumulants of arbitrary random variable

$$a = \sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}$$

in the graph W^* -probability space $(W^*(G), E)$.

1.1.1. Lattice Path Model.

Throughout this section, let G be a countable directed graph and let $(W^*(G), E)$ be the graph W^* -probability space over its diagonal subalgebra D_G . Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$ and let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in (W^*(G), E)$ be a D_G -valued random variable. In this section, we will define a lattice path model for the random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$. Recall that if $w = e_1 \dots e_k \in FP(G)$ with $e_1, \dots, e_k \in E(G)$, then we can define the length $|w|$ of w by k . i.e.e, the length $|w|$ of w is the cardinality k of the admissible edges $e_1, \dots, e_k$.

Definition 1.4. Let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semi-groupoid. If $w \in \mathbb{F}^+(G)$, then L_w is the corresponding D_G -valued random variable in $(W^*(G), E)$. We define the lattice path l_w of L_w and the lattice path l_w^{-1} of L_w^* by the lattice paths satisfying that ;

- (i) the lattice path l_w starts from $*$ $= (0, 0)$ on the $\mathbb{R}^2$ -plane.
- (ii) if $w \in V(G)$, then l_w has its end point $(0, 1)$.
- (iii) if $w \in E(G)$, then l_w has its end point $(1, 1)$.
- (iv) if $w \in E(G)$, then l_w^{-1} has its end point $(-1, -1)$.
- (v) if $w \in FP(G)$ with $|w| = k$, then l_w has its end point (k, k) .
- (vi) if $w \in FP(G)$ with $|w| = k$, then l_w^{-1} has its end point $(-k, -k)$.

Assume that finite paths $w_1, \dots, w_s$ in $FP(G)$ satisfy that $w_1 \dots w_s \in FP(G)$. Define the lattice path $l_{w_1 \dots w_s}$ by the connected lattice path of the lattice paths $l_{w_1}, \dots, l_{w_s}$. i.e.e, l_{w_2} starts from $(k_{w_1}, k_{w_1}) \in \mathbb{R}^+$ and ends at $(k_{w_1} + k_{w_2}, k_{w_1} + k_{w_2})$, where $|w_1| = k_{w_1}$ and $|w_2| = k_{w_2}$. Similarly, we can define the lattice path $l_{w_1 \dots w_s}^{-1}$ as the connected path of $l_{w_s}^{-1}, l_{w_{s-1}}^{-1}, \dots, l_{w_1}^{-1}$.

Definition 1.5. Let G be a countable directed graph and assume that $L_{w_1}, \dots, L_{w_n}$ are generators of $(W^*(G), E)$. Then we have the lattice paths $l_{w_1}, \dots, l_{w_n}$ of $L_{w_1}, \dots, L_{w_n}$, respectively in $\mathbb{R}^2$. Suppose that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in $(W^*(G), E)$, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Define the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of nonzero $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ by the connected lattice path of $l_{w_1}^{t_{w_1}}, \dots, l_{w_n}^{t_{w_n}}$, where $t_{w_j} = 1$ if $u_{w_j} = 1$ and $t_{w_j} = -1$ if

$u_{w_j} = *$. Assume that $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} = 0_{D_G}$. Then the empty set $\emptyset$ in $\mathbb{R}^2$ is the lattice path of it. We call it the empty lattice path. By LP_n , we will denote the set of all lattice paths of the D_G -valued random variables having their forms of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$, including empty lattice path.

Also, we will define the following important property on the set of all lattice paths ;

Definition 1.6. Let $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \neq \emptyset$ be a lattice path of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \neq 0_{D_G}$ in LP_n . If the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ starts from $*$ and ends on the $*$ -axis in $\mathbb{R}^+$, then we say that the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property. By LP_n^* , we will denote the set of all lattice paths having their forms of $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ which have the $*$ -axis-property. By little abuse of notation, sometimes, we will say that the D_G -valued random variable $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ satisfies the $*$ -axis-property if the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of it has the $*$ -axis-property.

The following theorem shows that finding $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}})$ is checking the $*$ -axis-property of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$.

Theorem 1.1. (See [15]) Let $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \in (W^*(G), E)$ be a D_G -valued random variable, where $u_{w_1}, \dots, u_{w_n} \in \{1, *\}$. Then $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) \neq 0_{D_G}$ if and only if $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ has the $*$ -axis-property (i.e., the corresponding lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ of $L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}$ is contained in LP_n^* . Notice that $\emptyset \notin LP_n^*$.) $\square$

By the previous theorem, we can conclude that $E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}) = L_v$, for some $v \in V(G)$ if and only if the lattice path $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}}$ has the $*$ -axis-property (i.e., $l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*$).

1.1.2. D_G -Valued Moments and Cumulants of Random Variables. Let $w_1, \dots, w_n \in \mathbb{F}^+(G)$, $u_1, \dots, u_n \in \{1, *\}$ and let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in (W^*(G), E)$ be a D_G -valued random variable. Recall that, in the previous section, we observed that the D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v \in (W^*(G), E)$ with $v \in V(G)$ if and only if the lattice path $l_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ of $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ has the $*$ -axis-property (equivalently, $l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*$). Throughout this section, fix a D_G -valued random variable $a \in (W^*(G), E)$. Then the D_G -valued random variable a has the following Fourier expansion,

$$a = \sum_{v \in V(G):a} p_v L_v + \sum_{l \in F P_*(G:a)} (p_l L_l + p_{l^t} L_{l^t}) + \sum_{w \in F P_*^c(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

Let's observe the new D_G -valued random variable $d_1 a d_2 a \dots d_n a \in (W^*(G), E)$, where $d_1, \dots, d_n \in D_G$ and $a \in W^*(G)$ is given. Put

$$d_j = \sum_{v_j \in V(G:d_j)} q_{v_j} L_{v_j} \in D_G, \text{ for } j = 1, \dots, n.$$

Notice that $V(G : d_j) = \mathbb{F}^+(G : d_j)$, since $d_j \in D_G \hookrightarrow W^*(G)$. Then

Proposition 1.2. (See [16]) *Let $a \in (W^*(G), E)$ be given as above. Then the n -th moment of a is*

$$E(d_1 a \dots d_n a) = \sum_{(v_1, \dots, v_n) \in \prod_{j=1}^n V(G:d_j)} (\prod_{j=1}^n q_{v_j})$$

$$\sum_{(w_1, \dots, w_n) \in \mathbb{F}^+(G:a)^n, u_{w_j} \in \{1, *\}, l_{w_1}^{u_{w_1}}, \dots, l_{w_n}^{u_{w_n}} \in LP_n^*} (\prod_{j=1}^n p_{w_j})$$

$$\left(\prod_{j=1}^n \delta_{(v_j, x_j, y_j; u_{w_j})} \right) E(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}}),$$

where

$$\delta_{(v_j, x_j, y_j)} = \begin{cases} \delta_{v_j, x_j} & \text{if } u_j = 1 \\ \delta_{v_j, y_j} & \text{if } u_j = * \end{cases}$$

□

Let $w_1, \dots, w_n \in FP(G)$ be finite paths and $u_1, \dots, u_n \in \{1, *\}$. Then, by the Möbius inversion, we have

$$(1.13) \quad k_n(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n}) = \sum_{\pi \in NC(n)} \widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \mu(\pi, 1_n),$$

where $\widehat{E} = (E^{(n)})_{n=1}^\infty$ is the moment multiplicative bimodule map induced by the conditional expectation E (See [16]) and where $NC(n)$ is the collection of all noncrossing partition over $\{1, \dots, n\}$.

Definition 1.7. *Let $NC(n)$ be the set of all noncrossing partition over $\{1, \dots, n\}$ and let $L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \in (W^*(G), E)$ be D_G -valued random variables, where $u_1, \dots, u_n \in \{1, *\}$. We say that the D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ is π -connected if the π -dependent D_G -moment of it is nonvanishing, for $\pi \in NC(n)$. In other words, the random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ is π -connected, for $\pi \in NC(n)$, if*

$$\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) \neq 0_{D_G}.$$

i.e., there exists a vertex $v \in V(G)$ such that

$$\widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}) = L_v.$$

For convenience, we will define the following subset of $NC(n)$;

Definition 1.8. Let $NC(n)$ be the set of all noncrossing partitions over $\{1, \dots, n\}$ and fix a D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ in $(W^*(G), E)$, where $u_1, \dots, u_n \in \{1, *\}$. For the fixed D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$, define

$$C_{w_1, \dots, w_n}^{u_1, \dots, u_n} \stackrel{\text{def}}{=} \{\pi \in NC(n) : L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \text{ is } \pi\text{-connected}\},$$

in $NC(n)$. Let μ be the Möbius function in the incidence algebra I_2 . Define the number $\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n}$, for the fixed D_G -valued random variable $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$, by

$$\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \stackrel{\text{def}}{=} \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \mu(\pi, 1_n).$$

Assume that there exists $\pi \in NC(n)$ such that $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$ is π -connected. Then $\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ and there exists the maximal partition $\pi_0 \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ such that $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$ is π_0 -connected. Notice that $1_n \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$. Therefore, the maximal partition in $C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ is 1_n . Hence we have that ;

Lemma 1.3. (See [16]) Let $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \in (W^*(G), E)$ be a D_G -valued random variable having the $*$ -axis-property. Then

$$E(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) = \widehat{E}(\pi)(L_{w_1}^{u_1} \otimes \dots \otimes L_{w_n}^{u_n}),$$

for all $\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}$. $\square$

By the previous lemmas, we have that

Theorem 1.4. (See [16]) Let $n \in 2\mathbb{N}$ and let $L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n} \in (W^*(G), E)$ be D_G -valued random variables, where $w_1, \dots, w_n \in FP(G)$ and $u_j \in \{1, *\}$, $j = 1, \dots, n$. Then

$$k_n(L_{w_1}^{u_1} \dots L_{w_n}^{u_n}) = \mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} \cdot E(L_{w_1}^{u_1}, \dots, L_{w_n}^{u_n}),$$

where $\mu_{w_1, \dots, w_n}^{u_1, \dots, u_n} = \sum_{\pi \in C_{w_1, \dots, w_n}^{u_1, \dots, u_n}} \mu(\pi, 1_n)$. $\square$

1.2. D_G -Freeness on $(W^*(G), E)$.

Now, we will introduce the diagram-distinctness of finite paths ;

Definition 1.9. (Diagram-Distinctness) We will say that the finite paths w_1 and w_2 are **diagram-distinct** if w_1 and w_2 have different diagrams in the graph G . Let X_1 and X_2 be subsets of $FP(G)$. The subsets X_1 and X_2 are said to be diagram-distinct if x_1 and x_2 are diagram-distinct for all pairs $(x_1, x_2) \in X_1 \times X_2$.

In [16], we found the D_G -freeness characterization on the generator set of $W^*(G)$, as follows ;

Theorem 1.5. (See [16]) Let $w_1, w_2 \in FP(G)$ be finite paths. The D_G -valued random variables L_{w_1} and L_{w_2} in $(W^*(G), E)$ are free over D_G if and only if w_1 and w_2 are diagram-distinct. $\square$

Let a and b be the given D_G -valued random variables. We can get the necessary condition for the D_G -freeness of a and b , in terms of their supports. Recall that we say that the two subsets X_1 and X_2 of $FP(G)$ are said to be diagram-distinct if x_1 and x_2 are diagram-distinct, for all pairs $(x_1, x_2) \in X_1 \times X_2$.

Proposition 1.6. (See [16]) Let $a, b \in (W^*(G), E)$ be D_G -valued random variables with their supports $\mathbb{F}^+(G : a)$ and $\mathbb{F}^+(G : b)$. The D_G -valued random variables a and b are free over D_G in $(W^*(G), E)$ if $FP(G : a_1)$ and $FP(G : a_2)$ are diagram-distinct. $\square$

2. DIAGONAL COMPRESSED RANDOM VARIABLES IN $(W^*(G), E)$

Let G be a countable directed graph and $(W^*(G), E)$, the graph W^* -probability space over the diagonal subalgebra D_G . In [18], we observed the diagonal compressed D_G -valued free probability on $(W^*(G), E)$. Fix a finite subset $V = \{v_1, \dots, v_N\}$ of the vertex set $V(G)$ and define the diagonal compression $P_V : W^*(G) \rightarrow W^*(G)$ by

$$P_V(a) = L_{v_1} a L_{v_1} + \dots + L_{v_N} a L_{v_N} \in W^*(G), \forall a \in (W^*(G), E).$$

If the given subset V is a singleton set, then we will call this diagonal compression the vertex-compression. Let $a \in (W^*(G), E)$ be a D_G -valued random variable having its expression

$$a = \sum_{v \in V(G:a)} p_v L_v + \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^t} L_l^*) + \sum_{w \in FP_*^c(G:a), u_w \in \{1, *\}} p_w L_w^{u_w}.$$

Then we have the following Fourier-like expression of the diagonal compressed random variable $P_V(a)$ of a by V ;

$$P_V(a) = \sum_{v \in V(G:P_V(a))} p_v L_v + \sum_{w \in FP_*(G:P_V(a))} (p_w L_w + p_{w^t} L_w^*) + \sum_{w \in FP_*^c(G:P_V(a)), u_w \in \{1, *\}} p_w L_w^{u_w},$$

with

$$V(G : P_V(a)) = V \cap V(G : a),$$

$$FP_*(G : P_V(a)) = (\cup_{j=1}^N \text{loop}_{v_j}(G : a)) \cap FP_*(G : a)$$

and

$$FP_*^c(G : P_V(a)) = (\cup_{j=1}^N \text{loop}_{v_j}(G : a)) \cap FP_*^c(G : a),$$

where

$$\text{loop}_{v_j}(G : a) = \{l \in \text{loop}(G : a) : l = v_j l v_j\},$$

for $j = 1, \dots, N$.

Hence we can apply all amalgamated free probability information on this diagonal compressed case. In particular, we can compute the D_G -valued moments and D_G -valued cumulants of the diagonal compressed random variables like Section 1.3 and hence we can get D_G -valued moment series and D_G -valued R-transforms of the diagonal compressed random variables (See [18]). Also, by little modification of Section 1.5, we can get the following theorems ;

Theorem 2.1. (See [18]) *Let $a, b \in (W^*(G), E)$ be a D_G -valued random variable and let $V = \{v_1, \dots, v_N\}$ be a finite subset of the vertex set $V(G)$. Let $P_V(a)$ and $P_V(b)$ be the diagonal compressed random variable of a and b by V in $(W^*(G), E)$, respectively. Then*

(1) *If a and b are free over D_G in $(W^*(G), E)$, then $P_V(a)$ and $P_V(b)$ are free over D_G in $(W^*(G), E)$.*

(2) *If a and b satisfy that*

$$(V \cap V(G : a)) \cap (V \cap V(G : b)) = \emptyset$$

and

$$\text{loop}_{v_i}(G : a) \cap \text{loop}_{v_j}(G : b) = \emptyset,$$

for all choices $(i, j) \in \{1, \dots, N\}^2$, then $P_V(a)$ and $P_V(b)$ are free over D_G in $(W^(G), E)$. $\square$*

Theorem 2.2. (See [18]) Let $a \in (W^*(G), E)$ be a D_G -valued random variable and let $V_1 = \{v_1^{(1)}, \dots, v_{N_1}^{(1)}\}$ and $V_2 = \{v_1^{(2)}, \dots, v_{N_2}^{(2)}\}$ be finite subsets of the vertex set $V(G)$. Suppose that

$$\text{loop}_{v_i^{(1)}}(G : a) \cap \text{loop}_{v_j^{(2)}}(G : a) = \emptyset,$$

for all choices $(i, j) \in \{1, \dots, N_1\} \times \{1, \dots, N_2\}$. Then the corresponding diagonal compressed random variables $P_{V_1}(a)$ and $P_{V_2}(a)$ of a by V_1 and V_2 are free over D_G in $(W^*(G), E)$. $\square$

Therefore, we can again characterize the D_G -freeness of the diagonal compressed random variables in $(W^*(G), E)$ by the subsets of the free semigroupoid $\mathbb{F}^+(G)$ of the graph G . Also, we can get the diagonal compressed R-transform calculus like in Section 1.6.

3. OFF-DIAGONAL RANDOM VARIABLES IN $(W^*(G), E)$

Throughout this chapter, let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semigroupoid of G and let $(W^*(G), E)$ be the graph W^* -probability space over the diagonal subalgebra D_G . In this chapter, we will consider the off-diagonal compressed random variables in $(W^*(G), E)$ over D_G . Let's fix $v_1 \neq v_2$ in $V(G)$ and let L_{v_1} and L_{v_2} be the corresponding projections in $(W^*(G), E)$. Define a subset $FP_{v_1}^{v_2}(G)$ of $FP(G)$ by

$$FP_{v_1}^{v_2}(G) \stackrel{\text{def}}{=} \{w \in FP(G) : w = v_1 w v_2\},$$

for $v_1, v_2 \in V(G)$.

Definition 3.1. Let $v_1 \neq v_2 \in V(G)$ be given. For any D_G -valued random variable $a \in (W^*(G), E)$, define the (v_1, v_2) -off-diagonal compressed random variable of a (in short (v_1, v_2) -compressed random variable of a) $_{v_1} a_{v_2}$ by

$$_{v_1} a_{v_2} \stackrel{\text{def}}{=} L_{v_1} a L_{v_2} \in (W^*(G), E).$$

Let $a \in (W^*(G), E)$ be an arbitrary D_G -valued random variable having the following Fourier expansion,

$$a = \sum_{v \in V(G:a)} p_v L_v + \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^*} L_l^*) + \sum_{w \in FP_*^c(G:a), u \in \{1, *\}} p_w L_w^u.$$

We will denote

$$a_d = \sum_{v \in V(G:a)} p_v L_v, \quad a_{(*)} = \sum_{l \in FP_*(G:a)} (p_l L_l + p_{l^t} L_l^*)$$

and

$$a_{(non-*)} = \sum_{w \in FP_*^c(G:a), u \in \{1, *\}} p_w L_w^u,$$

for the given D_G -valued random variable a . Thus the (v_1, v_2) -compressed random variable ${}_{v_1}a_{v_2}$ is determined by

$$\begin{aligned} {}_{v_1}a_{v_2} &= L_{v_1} a L_{v_2} = L_{v_1} a_d L_{v_2} + L_{v_1} a_{(*)} L_{v_2} + L_{v_1} a_{(non-*)} L_{v_2} \\ &= 0_{D_G} + L_{v_1} a_{(*)} L_{v_2} + L_{v_1} a_{(non-*)} L_{v_2}. \end{aligned}$$

By definition, we have the following partition of $FP(G : a)$, for the given random variable $a \in (W^*(G), E)$;

$$\{l = v_1 l v_2 : l \in FP_*(G : a)\} = FP_*(G : a) \cap FP_{v_1}^{v_2}(G)$$

and

$$\{w = v_1 w v_2 : w \in FP_*^c(G : a)\} = FP_*^c(G : a) \cap FP_{v_1}^{v_2}(G).$$

So, we have that ;

Lemma 3.1. *Let $a \in (W^*(G), E)$ be a D_G -valued random variable and let $v_1 \neq v_2 \in V(G)$ be the fixed vertices. Let ${}_{v_2}a_{v_1} \in (W^*(G), E)$ be the (v_1, v_2) -off-diagonal compressed random variable. Then*

$${}_{v_1}a_{v_2} = \sum_{w=v_1 w v_2 \in FP(G:a)} p_w L_w + \sum_{w'=v_2 w' v_1 \in FP(G:a)} p_{w'} L_{w'}^*.$$

Proof. By the relation that

$$L_w = L_{vw} = L_v L_w, \quad L_w = L_{wv'} = L_w L_{v'}$$

and

$$L_w^* = L_{wv}^* = L_v L_w^*, \quad L_w^* = L_{v'w}^* = L_w^* L_{v'},$$

under the weak topology $(v, v' \in V(G))$, we have that

$$L_w = L_{v w v'} = L_v L_w L_{v'}$$

and

$$L_w^* = L_{v w v'}^* = L_{v'} L_w^* L_v,$$

whenever $w = v w v'$ is a non-loop finite path, for $v, v' \in V(G)$. Thus, if $w \in FP(G : a)$, then we have that

$$L_{v_1} L_w L_{v_2} = \begin{cases} L_w & \text{if } w = v_1 w v_2 \\ 0_{D_G} & \text{otherwise} \end{cases}$$

and

$$L_{v_1} L_w^* L_{v_2} = \begin{cases} L_w^* & \text{if } w = v_2 w v_1 \\ 0_{D_G} & \text{otherwise.} \end{cases}$$

Therefore, the (v_1, v_2) -compressed random variable $v_1 a_{v_2}$ can be

$$\begin{aligned} v_1 a_{v_2} &= L_{v_1} a L_{v_2} = L_{v_1} \left(\sum_{w \in \mathbb{F}^+(G:a), u_w \in \{1,*\}} p_w L_w^{u_w} \right) L_{v_2} \\ &= L_{v_1} \left(\sum_{w \in FP(G:a), u_w \in \{1,*\}} p_w L_w^{u_w} \right) L_{v_2} \\ &= \sum_{w \in FP(G:a), u_w \in \{1,*\}} p_w (L_{v_1} L_w^{u_w} L_{v_2}) \\ &= \sum_{w=v_1 w v_2 \in FP(G:a)} p_w L_w + \sum_{w'=v_2 w' v_1 \in FP(G:a)} p_{w'} L_{w'}^*. \end{aligned}$$

■

For the convenience, we introduce the following new notation ;

Notation Let $v_1 \neq v_2 \in V(G)$ be given as before and let $a \in (W^*(G), E)$ be a D_G -valued random variable. Define

$$FP_{v_1}^{v_2}(G : a)_* \stackrel{def}{=} FP_*(G : a) \cap FP_{v_1}^{v_2}(G)$$

and

$$FP_{v_2}^{v_1}(G : a)_*^c \stackrel{def}{=} FP_*^c(G : a) \cap FP_{v_2}^{v_1}(G).$$

□

3.1. Off-Diagonal Compressed Moments and Cumulants.

Let $v_1 \neq v_2 \in V(G)$ be the fixed vertices. In this section, we will consider the moments and cumulants of the (v_1, v_2) -off-diagonal compressed random variables in the graph W^* -probability space $(W^*(G), E)$, over the diagonal subalgebra D_G . We have that

$$v_1 a_{v_2} = \sum_{w=v_1 w v_2 \in FP_{v_1}^{v_2}(G:a)} p_w L_w + \sum_{w'=v_2 w' v_1 \in FP_{v_2}^{v_1}(G:a)} p_{w'} L_{w'}^*.$$

So, to compute the D_G -valued moments and the D_G -valued cumulants of (v_1, v_2) -off-diagonal compressed random variables in $(W^*(G), E)$ is to compute the D_G -valued moments and the D_G -valued cumulants of the D_G -valued random variables $x \in (W^*(G), E)$ such that

$$x = \sum_{l_1=v_1 l_1 v_2 \in FP(G:x)} p_{l_1} L_{l_1} + \sum_{l_2=v_2 l_2 v_1 \in FP(G:x)} p_{l_2} L_{l_2}^*$$

in $(W^*(G), E)$.

Suppose that $a \in (W^*(G), E)$ is a D_G -valued random variable and assume that

$$FP_*(G : a) = \{w_1, w_2, \dots\} \subset FP(G).$$

Then, in terms of $FP_*(G : a)$, the D_G -valued random variable a has the following summands

$$p_{w_1} L_{w_1}, \quad p_{w_1^t} L_{w_1}^*, \quad p_{w_2} L_{w_2}, \quad p_{w_2^t} L_{w_2}^*, \dots,$$

where $p_{w_j}, p_{w_j^t} \in \mathbb{C}$. By the above observation, we have the following result ;

Theorem 3.2. *Let $v_1 \neq v_2 \in V(G)$ be the fixed vertices in the graph G and let $x \in (W^*(G), E)$ be a D_G -valued random variable with its Fourier-like expression,*

$$\sum_{l_1=v_1 l_1 v_2 \in FP(G:x)} p_{l_1} L_{l_1} + \sum_{l_2=v_2 l_2 v_1 \in FP(G:x)} p_{l_2} L_{l_2}^*.$$

Then the n -th moments and n -th cumulants of x vanish, for all $n \in \mathbb{N}$.

Proof. (1) Let $n = 1$. Then the first moments and the first cumulants of the D_G -valued random variable x vanish ;

$$E(x) = k_1(x) = 0_{D_G},$$

since $V(G : x) = \emptyset$.

(2) Let $n > 1$ in $\mathbb{N}$. Then the n -th D_G -valued moments vanish ; By Section 1.4, we have that the n -th moment of the D_G -valued random variable x is

$$\begin{aligned} & E(d_1 x \dots d_n x) \\ &= \sum_{\pi \in NC(n)} \sum_{(v^{(1)}, \dots, v^{(n)}) \in \prod_{j=1}^n V(G:d_j)} \left(\prod_{j=1}^n q_{v^{(j)}} \right) \end{aligned}$$

$$\sum_{(w_1, \dots, w_n) \in FP(G:x)^n, w_j = x_j w_j y_j, u_{w_j} \in \{1, *\}, l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*} \left(\prod_{j=1}^n p_{w_j} \right) \left(\prod_{j=1}^n \delta_{(v^{(j)}, x_j, y_j: u_{w_j})} \right) E \left(L_{w_1}^{u_1} \dots L_{w_n}^{u_n} \right),$$

where $x_j, y_j \in \{v_1, v_2\}$ and where $d_j = \sum_{v^{(j)} \in V(G: d_j)} q_{v^{(j)}} L_{v^{(j)}} \in D_G$ are arbitrary, $j = 1, \dots, n$, for all $n \in \mathbb{N}$, and where LP_n^* is the lattice path model satisfying the $*$ -axis-property (See Section 1.2). But to get the nonvanishing n -th cumulant of x , we need to have at least one summand of $d_1 a \dots d_n a$, $L_{w_1}^{u_1} \dots L_{w_n}^{u_n} = L_v$, for some $v \in V(G)$. Equivalently, the lattice path $l_{w_1, \dots, w_n}^{u_1, \dots, u_n}$ should have the $*$ -axis-property (i.e., $l_{w_1, \dots, w_n}^{u_1, \dots, u_n} \in LP_n^*$). To do that, at least, we need to have the nonempty $FP_*(G : x)$. But

$$V(G) \not\subseteq F^+(G : a) \text{ and } FP_*(G : x) = \emptyset.$$

Therefore,

$$k_1(d_1 x) = 0_{D_G} = E(d_1 a)$$

and

$$k_n(a, \dots, a) = k_n(a_{(*)}, \dots, a_{(*)}) = 0_{D_G}.$$

Indeed, we have that

$$FP(G : x) \subseteq FP_{v_1}^{v_2}(G) \cup FP_{v_2}^{v_1}(G).$$

Moreover, if $w = v_1 w v_2 \in FP(G : x) \cap FP_{v_1}^{v_2}(G)$, then the L_w -term of x exists but the L_w^* -term does not exist in the Fourier expansion of x . Similarly, if $w' = v_2 w v_1 \in FP(G : x) \cap FP_{v_2}^{v_1}(G)$, then the $L_{w'}^*$ -term of x exists but the $L_{w'}$ -term does not exist in the Fourier expansion of x . Therefore, each lattice path of $L_{w_1}^{u_1} \dots L_{w_n}^{u_n}$ does not have the $*$ -axis-property. Since all n -th D_G -valued cumulants of x vanish, all k -th D_G -valued moments of x vanish, by the Möbius inversion. ■

Consider the D_G -valued random variable

$$a = L_v + L_{w_1} + L_{w_1}^* + L_{w_2}^*,$$

where $v \in V(G)$ and $w_1 \neq w_2 \in FP(G)$, with $w_1 = v_1 w_1 v_2$ and $w_2 = v_2 w_2 v_1$. Then the (v_1, v_2) -off-diagonal compressed random variable of the D_G -valued random variable a is

$$_{v_1} a_{v_2} = L_{w_1} + L_{w_2}^*.$$

So, we have that

$$FP_*(G : _{v_1} a_{v_2}) = \emptyset$$

and

$$FP_*^c(G : {}_{v_1}a_{v_2}) = \{w_1, w_2\}.$$

Therefore, the n -th moments and the n -th cumulants of ${}_{v_1}a_{v_2}$ vanish, for all $n \in \mathbb{N}$.

3.2. Off-Diagonal Compressed D_G -Freeness.

Let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semigroupoid of G and let $(W^*(G), E)$ be the corresponding graph W^* -probability space over the diagonal subalgebra D_G . In this section, we will consider the D_G -freeness of the off-diagonal compressed random variables. Throughout this section, let v_1, v_2, v_3 and v_4 be mutually distinct vertices in $V(G)$. Since $FP_*(G : {}_{v_1}a_{v_2}) = \emptyset$, it has vanishing D_G -valued n -th moments and n -th cumulants, for all $n \in \mathbb{N}$. Therefore, automatically, the D_G -valued moments vanish, by [12].)

Proposition 3.3. *Let a and b be D_G -valued random variables in the graph W^* -probability space $(W^*(G), E)$ and let $v_1 \neq v_2$ be the given vertices in $V(G)$. If a and b have the diagram-distinct supports, then the (v_1, v_2) -off-diagonal compressed random variables ${}_{v_1}a_{v_2} \equiv L_{v_1}aL_{v_2}$ and ${}_{v_1}b_{v_2} \equiv L_{v_1}bL_{v_2}$ are free over D_G in $(W^*(G), E)$.*

Proof. By the diagram-distinctness of $FP(G : a)$ and $FP(G : b)$, $loop^c(G : a)$ and $loop^c(G : b)$ are diagram-distinct, too. Note that

$$\begin{aligned} & FP(G : {}_{v_1}a_{v_2}) \subset loop^c(G : a) \\ \text{and} \\ & FP(G : {}_{v_1}b_{v_2}) \subset loop^c(G : b). \end{aligned}$$

Therefore, ${}_{v_1}a_{v_2}$ and ${}_{v_1}b_{v_2}$ are free over D_G . ■

Now, we will consider the other case ;

Proposition 3.4. *Let v_1, v_2, v_3 and v_4 be the mutually distinct vertices in $V(G)$ and let $a \in (W^*(G), E)$ be a D_G -valued random variable. Then the (v_1, v_2) -off-diagonal compressed random variable ${}_{v_1}a_{v_2}$ and the (v_3, v_4) -off-diagonal compressed random variable ${}_{v_3}a_{v_4}$ are free over D_G in $(W^*(G), E)$.*

Proof. Since $v_1, \dots, v_4$ are mutually distinct, the (v_1, v_2) -off diagonal compressed random variable ${}_{v_1}a_{v_2}$ and the (v_3, v_4) -off-diagonal compressed random variable ${}_{v_3}a_{v_4}$ have the diagram-distinct supports. Therefore, they are free over D_G in $(W^*(G), E)$. ■

Corollary 3.5. *Let v_1, v_2, v_3 and v_4 be the given vertices in $V(G)$. Define two subsets*

$$W^*(G)_{v_1}^{v_2} = L_{v_1} W^*(G) L_{v_2} \quad \text{and} \quad W^*(G)_{v_3}^{v_4} = L_{v_3} W^*(G) L_{v_4},$$

in the graph W^ -probability space $(W^*(G), E)$. Then these subsets are free over D_G in $(W^*(G), E)$. $\square$*

We can regard $W^*(G)_{v_i}^{v_j}$ as an off-diagonal block of $W^*(G)$.

4. COMPRESSED FREE PROBABILITY ON $(W^*(G), E)$

Throughout this chapter, let G be a countable directed graph and $\mathbb{F}^+(G)$, the free semigroupoid of the graph G and let $(W^*(G), E)$ be the graph W^* -probability space over the diagonal subalgebra D_G . In this chapter, we will consider the compressed random variable PaP of the D_G -valued random variable $a \in (W^*(G), E)$ by the projection $P \in W^*(G)$. Let $v_1, \dots, v_N$ be vertices in $V(G)$ and define the projection

$$P = L_{v_1} + \dots + L_{v_N} \in W^*(G).$$

Then it is indeed a projection in $W^*(G)$. From now, fix the finite vertices $v_1, \dots, v_N \in V(G)$ and the corresponding projection $P = L_{v_1} + \dots + L_{v_N}$ in $W^*(G)$. Let $a \in (W^*(G), E)$ be a D_G -valued random variable. Then naturally, we can construct the compressed random variable PaP of a by P . Then this compressed random variable is again a D_G -valued random variable in $(W^*(G), E)$.

Notice that the diagonal-compressed random variable $P_V(a)$ by the diagonal compression $P_V : W^*(G) \rightarrow W^*(G)$, for the fixed vertex-subset $V = \{v_1, \dots, v_N\}$ (See Chapter 2 and [18]) and the compressed random variable PaP are totally different in $(W^*(G), E)$. For example, if a is a D_G -valued random variable in $(W^*(G), E)$, then the diagonal compressed random variable is

$$P_V(a) \stackrel{def}{=} L_{v_1} a L_{v_1} + \dots + L_{v_N} a L_{v_N}$$

but the compressed random variable by the projection P is

$$PaP = \left(\sum_{i=1}^N L_{v_i} \right) a \left(\sum_{j=1}^N L_{v_j} \right) = \sum_{(i,j) \in \{1, \dots, N\}^2} L_{v_i} a L_{v_j}.$$

Therefore, we can say that the compressed random variable PaP of a by P satisfies that

$$PaP = P_V(a) + \sum_{(i,j) \in \{1, \dots, N\}^2, i \neq j} L_{v_i} a L_{v_j}.$$

However, in this chapter, we will observe that $P_V(a)$ and PaP have the same free probability information.

Again, remark that the compressed random variable PaP is the sum of diagonal compressed random variable $P_V(a)$ and the (sum of D_G -free) off-diagonal compressed random variables $L_{v_i} a L_{v_j}$ ($i \neq j$ in $\{1, \dots, N\}$). Define

$$P_V^c(a) = \sum_{(i,j) \in \{1, \dots, N\}^2, i \neq j} L_{v_i} a L_{v_j}.$$

Proposition 4.1. *Let $a \in (W^*(G), E)$ be a D_G -valued random variable and let $P = \sum_{j=1}^N L_{v_j} \in W^*(G)$ be a projection, where $V = \{v_1, \dots, v_N\} \subset V(G)$ is the finite subset of $V(G)$. Then the compressed random variable PaP of a by P is*

$$PaP = P_V(a) + P_V^c(a),$$

where $P_V(a)$ is the diagonal compressed random variable of a by V and

$$P_V^c(a) = \sum_{(i,j) \in \{1, \dots, N\}^2, i \neq j} L_{v_i} a L_{v_j},$$

In particular, $P_V(a)$ and $P_V^c(a)$ are free over D_G in $(W^*(G), E)$.

Proof. By the previous discussion, the compressed random variable of a by the projection P satisfies that

$$PaP = P_V(a) + P_V^c(a),$$

where

$$P_V(a) = L_{v_1} a L_{v_1} + \dots + L_{v_N} a L_{v_N}$$

and

$$P_V^c(a) = \sum_{(i,j) \in \{1, \dots, N\}^2, i \neq j} L_{v_i} a L_{v_j}.$$

Then, by [18] and by Chapter 3,

$$\mathbb{F}^+(G : P_V(a)) \subset (V \cup \text{loop}(G))$$

and

$$\mathbb{F}^+(G : P_V^c(a)) \subset \text{loop}^c(G).$$

Therefore, the supports of $P_V(a)$ and $P_V^c(a)$ are diagram-distinct and hence they are free over D_G . ■

4.1. Amalgamated Moments and Cumulants of Compressed Random Variables.

Remark that the compressed random variable PaP of a D_G -valued random variable $a \in (W^*(G), E)$ by the projection $P = \sum_{j=1}^N L_{v_j} \in W^*(G)$ has the form of

$$PaP = P_V(a) + P_V^c(a),$$

where $V = \{v_1, \dots, v_N\} \subset V(G)$ is the finite subset. Furthermore, by the previous proposition, as D_G -valued random variables in $(W^*(G), E)$, the diagonal compressed part $P_V(a)$ and the off-diagonal compressed part $P_V^c(a)$ are free over D_G in $(W^*(G), E)$. Therefore, we can get the following result ;

Theorem 4.2. *Let $V = \{v_1, \dots, v_N\}$ be the finite subset of the vertex set $V(G)$ and let $P = \sum_{j=1}^N L_{v_j} \in W^*(G)$ be the corresponding projection. Let $a \in (W^*(G), E)$ be a D_G -valued random variable and PaP , the compressed random variable of a by P . Then the n -th cumulants of PaP is*

$$k_1(d_1 PaP) = \sum_{v \in V \cap (V(G:d_1) \cap V(G:a))} (q_v p_v) L_v$$

and

$$\begin{aligned} & k_n \left(\underbrace{d_1 PaP, \dots, d_n PaP}_{n\text{-times}} \right) \\ &= \sum_{(v^{(1)}, \dots, v^{(n)}) \in \Pi_{k=1}^n V(G:d_k)} \left(\prod_{j=1}^n q_{v^{(j)}} \right) \\ & \quad \sum_{(w_1, \dots, w_n) \in \left(\left(\bigcup_{k=1}^n \text{loop}_{v_k}(G:a) \right) \cup V \cap V(G:a) \right)^n, w_j = x_j w_j x_j, l_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} \in LP_n^*} \\ & \quad \left(\prod_{j=1}^n p_{w_j} \right) \left(\prod_{k=1}^n \delta_{v^{(k)}, x_k} \right) \mu_{w_1, \dots, w_n}^{u_{w_1}, \dots, u_{w_n}} E \left(L_{w_1}^{u_{w_1}} \dots L_{w_n}^{u_{w_n}} \right), \end{aligned}$$

for all $n > 1$ in $\mathbb{N}$, where $d_k = \sum_{v^{(k)} \in V(G:d_k)} q_{v^{(k)}} L_{v^{(k)}} \in D_G$ are arbitrary for $k = 1, \dots, n$.

Proof. Suppose we have the compressed random variable PaP of the D_G -valued random variable a . Then

$$PaP = P_V(a) + P_V^c(a),$$

where P_V is the diagonal compression and P_V^c is the off-diagonal compression by $V \subset V(G)$. Clearly, we have the above first cumulant of PaP . Also, we have that

$$\begin{aligned} & k_n(d_1 PaP, \dots, d_n PaP) \\ &= k_n(d_1(P_V(a) + P_V^c(a)) + \dots + d_n(P_V(a) + P_V^c(a))) \\ &= k_n(d_1 P_V(a), \dots, d_n P_V(a)) \\ &\quad + k_n(d_1 P_V^c(a), \dots, d_n P_V^c(a)) \end{aligned}$$

by the D_G -freeness of $P_V(a)$ and $P_V^c(a)$

$$\begin{aligned} &= k_n(d_1 P_V(a), \dots, d_n P_V(a)) \\ &\quad + \sum_{(i,j) \in \{1, \dots, N\}^2, i \neq j} k_n(d_1(L_{v_i} a L_{v_j}), \dots, d_n(L_{v_i} a L_{v_j})) \\ &= k_n(d_1 P_V(a), \dots, d_n P_V(a)) + 0_{D_G} \end{aligned}$$

by Section 3.1

$$= k_n(d_1 P_V(a), \dots, d_n P_V(a)),$$

for all $n \in 2\mathbb{N}$. Therefore, by [18], we can get the above result. ■

Remark 4.1. *The above theorem simply shows that*

$$k_n(d_1(PaP), \dots, d_n(PaP)) = k_n(d_1 P_V(a), \dots, d_n P_V(a)),$$

for all $n \in \mathbb{N}$ and for any arbitrary $d_1, \dots, d_n \in D_G$.

This says that the off-diagonal compressed part $P_V^c(a)$ does not affect to compute the D_G -valued cumulants of the compressed random variable PaP . We can conclude that the compressed random variable PaP of the D_G -valued random variable by $P = \sum_{j=1}^N L_{v_j}$ and the diagonal compressed random variable $P_V(a)$ of the random variable a by $V = \{v_1, \dots, v_N\}$ have the same distributions and hence they have the same D_G -valued R-transforms.

4.2. D_G -Freeness of Compressed Random Variables.

In this section, we will consider the D_G -freeness of compressed random variables. In this section, we will consider the various conditions for the D_G -freeness of compressed random variables.

Theorem 4.3. *Let $a, b \in (W^*(G), E)$ be a D_G -valued random variable and let $V = \{v_1, \dots, v_N\}$ be a finite subset of the vertex set $V(G)$ and $P = \sum_{j=1}^N L_{v_j}$, the corresponding projection in $W^*(G)$. Let PaP and PbP be the compressed random variable of a and b by P in $(W^*(G), E)$, respectively. If a and b satisfy that*

$$(V \cap V(G : a)) \cap (V \cap V(G : b)) = \emptyset,$$

$$\text{and} \quad \text{loop}^c(G : a) \cap \text{loop}^c(G : b) = \emptyset,$$

$$\text{loop}_{v_i}(G : a) \cap \text{loop}_{v_j}(G : b) = \emptyset,$$

for all choices $(i, j) \in \{1, \dots, N\}^2$, then PaP and PbP are free over D_G in $(W^(G), E)$.*

Proof. By the previous section, we have that

$$PaP = P_V(a) + P_V^c(a) \quad \text{and} \quad PbP = P_V(b) + P_V^c(b),$$

where $P_V, P_V^c : W^*(G) \rightarrow W^*(G)$ are the diagonal compression and off-diagonal compression by V , respectively. Moreover, $P_V(a)$ (resp. $P_V(b)$) and $P_V^c(a)$ (resp. $P_V^c(b)$) are free over D_G in $(W^*(G), E)$. By the assumption and by [18], $P_V(a)$ and $P_V(b)$ are free over D_G in $(W^*(G), E)$. By the second condition $P_V^c(a)$ and $P_V^c(b)$ are free over D_G in $(W^*(G), E)$. This shows that $\{P_V(a), P_V^c(a)\}$ and $\{P_V(b), P_V^c(b)\}$ are free over D_G in $(W^*(G), E)$. So, PaP and PbP are free over D_G . ■

Theorem 4.4. *Let $a \in (W^*(G), E)$ be a D_G -valued random variable and let $V_1 = \{v_1^{(1)}, \dots, v_{N_1}^{(1)}\}$ and $V_2 = \{v_1^{(2)}, \dots, v_{N_2}^{(2)}\}$ be finite subsets of the vertex set $V(G)$. Let P and Q be the corresponding projections of V_1 and V_2 in $W^*(G)$, respectively. Suppose that*

$$(V_1 \cap V(G : a)) \cap (V_2 \cap V(G : a)) = \emptyset,$$

$$\text{and} \quad \text{loop}_{v_i^{(1)}}(G : a) \cap \text{loop}_{v_j^{(2)}}(G : a) = \emptyset,$$

for all choices $(i, j) \in \{1, \dots, N_1\} \times \{1, \dots, N_2\}$. Then the corresponding diagonal compressed random variables PaP and QaQ of a by P and Q are free over D_G in $(W^(G), E)$.*

Proof. By hypothesis and by [18], $P_{V_1}(a)$ and $Q_{V_2}(a)$ are free over D_G in $(W^*(G), E)$, where $P_{V_1}, Q_{V_2} : W^*(G) \rightarrow W^*(G)$ are diagonal compressions by V_1 and V_2 , respectively. ■

Reference

- [1] A. Nica, R-transform in Free Probability, IHP course note, available at www.math.uwaterloo.ca/~anica.
- [2] A. Nica and R. Speicher, R-diagonal Pair-A Common Approach to Haar Unitaries and Circular Elements, (1995), www.mast.queensu.ca/~speicher.
- [3] B. Solel, You can see the arrows in a Quiver Operator Algebras, (2000), preprint
- [4] A. Nica, D. Shlyakhtenko and R. Speicher, R-cyclic Families of Matrices in Free Probability, J. of Funct Anal, 188 (2002), 227-271.
- [5] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.
- [6] D. Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).
- [7] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.
- [8] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.
- [9] D.W. Kribs and M.T. Jury, Ideal Structure in Free Semigroupoid Algebras from Directed Graphs, preprint
- [10] D.W. Kribs and S.C. Power, Free Semigroupoid Algebras, preprint
- [11] I. Cho, Amalgamated Boxed Convolution and Amalgamated R-transform Theory, (2002), preprint.
- [12] I. Cho, The Tower of Amalgamated Noncommutative Probability Spaces, (2002), Preprint.
- [13] I. Cho, Compatibility of a Noncommutative Probability Space and a Noncommutative Probability Space with Amalgamation, (2003), Preprint
- [14] I. Cho, An Example of Scalar-Valued Moments, Under Compatibility, (2003), Preprint.
- [15] I. Cho, Free Semigroupoid Probability Theory, (2004), Preprint.
- [16] I. Cho, Graph W^* -Probability Theory, (2004), Preprint.
- [17] I. Cho, Random Variables in Graph W^* -Probability Spaces, (2004), Preprint.
- [18] I. Cho, Diagonal Compressed Random Variables in the Graph W^* -Probability Space, (2004), Preprint.
- [19] I. Cho, Free Product of Two Graph W^* -Probability Spaces, (2004), Preprint.
- [20] P. Śniady and R. Speicher, Continuous Family of Invariant Subspaces for R-diagonal Operators, Invent Math, 146, (2001) 329-363.
- [21] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).

[22] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at www.mast.queensu.ca/~speicher.

UNIV. OF IOWA, DEP. OF MATH, IOWA CITY, IA, U. S. A.,
E-mail address: `ilcho@math.uiowa.edu`